

NATURAL RADIOACTIVITY IN MAJOR RIVERS OF KELANTAN STATE, MALAYSIA

Nuraddeen Nasirum Garba^{a,b*}, Ahmad Termizi Ramli^a, Muneer Aziz Saleh^{c,d}, Mohd Syazwan Mohd Sanusi^a, Hamman Tukur Gabdo^a, Abubakar Sadiq Aliyu^{a,e}

Article history

Received

15 April 2015

Received in revised form

29 September 2015

Accepted

12 November 2015

*Corresponding author
nurangetso@yahoo.com

^aDepartment of Physics, Faculty of Science, Universiti Teknologi Malaysia, 81310 Johor, Malaysia


^bDepartment of Physics, Ahmadu Bello University, Zaria, Nigeria

^cNuclear Engineering Programme, Faculty of Petroleum and Renewable Energy Engineering, Universiti Teknologi Malaysia, 81310 Johor, Malaysia

^dNational Atomic Energy Commission (NATEC), Sana'a, Yemen

^eDepartment of Physics, Nassarawa State University Keffi, Nigeria

Graphical abstract



Abstract

Assessment of natural radionuclides (^{238}U , ^{232}Th , and ^{40}K) and terrestrial gamma radiation dose rates (TGRD) in major rivers of Kelantan states, Malaysia were conducted. Measurements were carried out using a portable [NaI(Tl)] micro roentgen (μR) survey meter and inductively coupled plasma mass spectrometer (ICP-MS) for in situ TGRD and the activity concentrations of ^{238}U , ^{232}Th , and ^{40}K in water samples, respectively. The mean TGRD was found to be $312.98 \text{ nGy h}^{-1}$ and from water samples analyzed, the mean activity concentrations of ^{238}U , ^{232}Th , and ^{40}K was found to be 13.31 mBq L^{-1} , 4.39 mBq L^{-1} and $1118.72 \text{ mBq L}^{-1}$ which were about 3 times and slightly higher than the world average values of 5 mBq L^{-1} and 3 mBq L^{-1} respectively.

Keywords: Radioactivity, ICP-MS, Major Rivers, TGRD, AGDE

© 2015 Penerbit UTM Press. All rights reserved

1.0 INTRODUCTION

Water is the most abundant substance on earth and is the principal constituent of all living things. Man uses water for many purposes such as power generation, agriculture and above all for domestic purposes.

Water for human consumption should be free from chemical, microbiological and radiological contamination [1]. Radioactivity in terrestrial, fresh water and marine media arises mainly from the primordial radionuclides decay series of ^{238}U , ^{232}Th and ^{40}K . Radiological impact assessment to the population in both human and non-human biota system as a result of radiation emitted by these natural radionuclides is important, since they contribute to the collective dose of the population [2, 3]. Exposure to

ionizing radiation from natural sources is a continuous and unavoidable feature of life on earth [4].

In order to limit radiation exposure to members of the public as low as reasonably as achievable (ALARA), assessment of radionuclide concentrations in water has become necessary. [5]. Radiation exposure through water may be due to ingestion (drinking water) and inhalation (from household water), both of these mechanisms pose potential health hazard [6-8]. This study is the first of its kind, which aim to measure the natural radionuclides levels in major rivers of Kelantan State, Malaysia. The data obtained will be useful in assessing the radiological health implications associated with the natural radionuclides, which will be needed in formulating safety standards and national guidelines for Malaysia in light of international recommendations.

2.0 EXPERIMENTAL

2.1 Study Area

Kelantan state is located in the North-Eastern part of Peninsula Malaysia, covering latitude 4° 0' to 5° 15' N and longitude 101° 15' to 102° 45'. It is tropic with

temperature ranges from 21 to 32 °C and intermittent rain fall throughout the year. It is located on granites set which is undifferentiated intrusive rock [9, 10]. The state consists of eight underlying geological formations (Table 1) which was overlain by twelve soil types (Table 2).

Table 1 Geological formations of Kelantan [11, 12]

S/N	Name	Description
1	Quaternary (1)	Composed of continental and marine deposits with unconsolidated sand mainly marine.
2	Quaternary (2)	Composed of continental and marine deposits with unconsolidated silt and clay (main).
3	Quaternary (4)	Consists of marine and continental deposits with unconsolidated clay, silt, sand and gravel undifferentiated.
4	Triassic (14)	Characterized by lithologies consisting of interbedded sandstone, silt, shale, conglomerate and chert.
5	Permian (21)	Consists of shale, slate, and phyllite with subordinate schist and sandstone. It developed of limestone through the succession. Rhyolitic, volcanic to andesitic in composition widespread, with unconsolidated deposit of ignimbrite.
6	Permian (22)	Consists of Permian (21), with unconsolidated deposit of ignimbrite.
7	Silurian Ordovician (35)	Underlain by acid intrusion and overlain by Triassic which shows that the area has a contact with sea water being leached out under pressure.
8	Acid Intrusive (38)	Consists of undifferentiated igneous rocks; they are mostly granitic rocks.

Table 2 Soil types of Kelantan [12, 13]

Soil type label	FAO UNIT	Local Name
1	Humic Podzols -Dystric Fluvisols	Rudua- Rusila
10	Dystric Histosols	Gambut
11	Dystric Fluvisols - Dystric Gleysols	Telemong Akob- Tanah Lanar Tempatan
12	Haplic Acrisol- Haplic Acrisol-Gleyic Luvisol	Tokyong- Chempaka – lundang
13	Dystric Fluvisols- Gleyic Acrisol	Lubok itek- lubok sending
14	Gleyic Nitosols- Acric Ferralsols	Batu Hiram- Kampong Tepus- Sungai Amin
17	Xanthic Ferrasols -Dystric Gleysols	Holyrood Lunas
25	Plinthic Ferralsols- Plinthic Ferralsols- Plinthic ferralsol	Melaka- Tavy- Gajah Mati
32	Dystric Nitosols - Orthic Ferrasols	Rengam Jerangau
37	Ferric Acrisols- Orthic Ferralsols	Batang Merbau – Munchong
39	Ferric Acrisols- Orthic Ferralsols- Ferric Acrisols	Durian – Munchong – Bungor
4	Steep land	Tanah churam

2.2 Sample Collection

2 liters of water samples were collected each from major rivers in Kelantan. A total of five samples were collected and the coordinates of each of the

sampling locations were recorded with the aid of Global Positioning System, Garmin (GPS Map76). The samples were collected in an airtight plastic bottles with minimal aeration, and about 2 mL of nitric acid (HNO₃) were added to avoid ions from being

adsorbed to the bottle wall [14] and then transported to the laboratory for preparation and analysis.

2.3 Sample Preparation and Analysis

The samples were filtered through a Whatman no.42 filter paper and then 10 mL of each sample was placed in a test tube for analysis. The analysis was performed using an inductively coupled plasma mass spectrometer (ICP-MS) instrument model ELAN 6000 at the Malaysian Nuclear Agency. A blank solution was run to estimate the background activity of ^{238}U , ^{232}Th and ^{40}K due to the system. The measurement procedure includes use of calibration standard and two quality control (QC) samples for each set of the samples.

3.0 RESULTS AND DISCUSSION

A details TGRD assessment of Kelantan was conducted by Garba et al. [12], and reported a

mean TGRD values of 209 nGy h⁻¹ and 150 nGy h⁻¹, which was more than twice the world average value of 59 nGy h⁻¹. Due to the anomaly observed in the measured TGRD, it was expected that water resources of the area might be affected by this high background radioactivity usually caused by uranium and thorium.

The activity concentrations of ^{238}U , ^{232}Th and ^{40}K in the samples measured using ICP-MS technique and the measured TGRD at each sampling point were presented in Table 3. The mean activity concentration of ^{238}U , ^{232}Th and ^{40}K were found to be 13.31 mBq L⁻¹ and 1118.72 mBq L⁻¹, respectively. It can be seen from (Table 3) that the mean activity concentrations of ^{238}U and ^{232}Th (13.31 and 4.39) mBq L⁻¹ in Kelantan rivers are about 3 times and slightly higher than the world average values of 5 mBq L⁻¹ and 3 mBq L⁻¹ respectively [15], and are also about 15 and 88 times higher than the corresponding values reported for New York City which was 0.9 mBq L⁻¹ and 0.05 mBq L⁻¹ for ^{238}U and ^{232}Th respectively [16] and 3 and 2 times higher than that of north western

Table 3 Activity concentrations of ^{238}U , ^{232}Th and ^{40}K and measured TGRD rates

Location	Sample Id	D (nGy h ⁻¹)	^{40}K (mBq L ⁻¹)	^{238}U (mBq L ⁻¹)	^{232}Th (mBq L ⁻¹)	($^{232}\text{Th}/^{238}\text{U}$) Ratio
Hutan Rizab Gunung Sabab	KLTN W1	413.25	68.92 ± 0.91	0.79 ± 0.01	0.36 ± 0.04	0.46
Kolam Air Panas Batu Melati	KLTN W2	380.63	5259.85 ± 151.15	31.66 ± 5.42	0.44 ± 0.04	0.01
Kg. Pasar Era	KLTN W3	326.25	72.55 ± 0.06	10.68 ± 0.06	3.38 ± 0.01	0.32
Kg. Sg. Sok	KLTN W4	324.08	99.15 ± 0.06	14.17 ± 0.49	13.19 ± 0.04	0.93
Kg. Rela Jabo	KLTN W5	120.71	93.11 ± 0.30	9.25 ± 0.26	4.56 ± 0.24	0.49
Kelantan State	Mean	312.98	1118.72 ± 30.71	13.31 ± 1.25	4.39 ± 0.07	0.33 ~ 1:3

Pakistan with the values of 3.9 mBq L⁻¹ and 2 mBq L⁻¹ as well as about 1.5, 2 and 4 times higher than those reported for Pahang rivers, Malaysia which are 8.49 mBq L⁻¹, 1.74 mBq L⁻¹ and 77 mBq L⁻¹ for ^{238}U , ^{232}Th and ^{40}K respectively [17, 18].

This can be attributed to the geological formations of Kelantan which is predominantly granitic in nature. Granites and metamorphic areas are rich in uranium, thorium and potassium with uranium having higher transfer potential to water due to high solubility of some of its compound [19]. Previous studies in other areas shows high concentration of uranium, thorium and potassium in areas covered by granites and metamorphic rocks [20, 21]. The mean ratio of thorium to uranium (Th/U) concentration in the rivers were found to be 0.33 and 0.32 which stands for 1:3 for both rivers, and this is due to the higher solubility of uranium in water.

A graph showing the variation of ^{238}U , ^{232}Th and ^{40}K concentrations of in the major rivers was plotted and presented below as Figure 1.




Figure 1 Activity concentrations of ^{238}U , ^{232}Th and ^{40}K

High concentrations of ^{238}U and ^{40}K in Kelantan was observed in sample with Id KLTN W2 which was collected from Kolam Air Panas Batu Melati (Hot Spring) with a value of 5259.85 and 31.66 mBq L⁻¹ respectively, which flows and spread over a granitic

rock and serve as a tourist attraction site while high ^{232}Th concentration was observed in sample with Id KLTN W4 collected from Kg. Sg. Sok with a value of 13.19 mBq L^{-1} , which exceeded the world average values of 5 mBq L^{-1} and 3 mBq L^{-1} respectively [15], while low concentrations ^{238}U and ^{232}Th were observed in sample with Id KLTN W1 and that of ^{40}K in sample with Id KLTN W5 collected from Hutan Rizab Gunung Sabab and Kg. Rela Jabo respectively both in Kelantan with respective values of 0.79, 0.36 and $120.71 \text{ mBq L}^{-1}$.

However, the high concentrations of the radionuclides observed in some of the samples collected can be attributed to geological formations of the areas which mostly consists of granites and gneisses which contained high concentration of ^{238}U , ^{232}Th series and ^{40}K [5] while some may be due to continuous movement of water from nearby granitic region. Samples with Id's KLTN W2 and KLTN W4 were collected from areas underlain by acid intrusive and Triassic geological formations. These areas have high TGRD of values $380.63 \text{ nGy h}^{-1}$ and $324.08 \text{ nGy h}^{-1}$ respectively. On the other hand with the lowest value for both ^{238}U and ^{232}Th were from areas dominated by Carboniferous geological formation which was known to be originated from sedimentary rocks than that originated from igneous rock.

4.0 CONCLUSION

TGRD and ^{238}U , ^{232}Th and ^{40}K activity concentrations in major rivers of Kelantan was carried out. The mean TGRD was found to be $312.98 \text{ nGy h}^{-1}$ with specific activity concentrations of ^{238}U , ^{232}Th and ^{40}K found to be 13.31 mBq L^{-1} , 4.39 mBq L^{-1} and $1118.72 \text{ mBq L}^{-1}$ for Kelantan respectively. The mean activity concentrations of ^{238}U , ^{232}Th and ^{40}K obtained was found to be about 3 times and slightly higher than the world average values of 5 mBq L^{-1} and 3 mBq L^{-1} respectively, which can be attributed to the geological formations in the state which was predominantly granitic in nature.

Acknowledgement

This project is funded by the Atomic Licensing Board (AELB) of Malaysia, Ministry of Science, Technology and Innovation, Malaysia under project title: PEMETAAN ISODOS SINARAN GAMA DATARAN, SEMENANJUANG MALAYSIA, which involved Lee, M.H., Heryanshah, A., Wagiran, H., and Said, M.N. The consultancy was managed by GITM Sdn Bhd. The authors would like to thank the Ministry of Higher Education Malaysia (MOHE) and Universiti Teknologi Malaysia (UTM) for support and funding under UTM Research University Grant; QJ130000.2526.03H67.

References

- [1] Garba, N. N., N. Rabi and B. B. M. Dewu. 2012. Preliminary Studies on ^{222}Rn Concentration in Ground Water from Zaria Metropolis. *Journal of Physical Science*. 23: 1-9.
- [2] UNSCEAR. 1988. Sources, Effects and Risks of Ionizing Radiation. United Nations Scientific Committee on the Effects of Atomic Radiation, United Nations, New York.
- [3] UNSCEAR. 2008. Sources and Effects of Ionizing Radiation, in United Nation Publication, R. United Nations Scientific Committee on the Effects of Atomic, Editor: New York.
- [4] UNSCEAR. 2000. Effects of Ionizing Radiation, 2000 Report to the General Assembly, with Scientific Annexes. United Nations, New York.
- [5] Ahmad T. A., R. A. Termizi, and W. Abdul-Khalik. 2004. Analysis of the Concentration of Natural Radionuclides in Rivers in Kota Tinggi District, Malaysia. *Journal of Nuclear and Related Technologies*. 1: 41-52.
- [6] Savidou, A., G. Sideris, and N. Zouridakis. 2001. Radon in Public Water Supplies in Migdonia Basin, Central Macedonia, Northern Greece. *Health Physics*. 80: 170-174.
- [7] Sohrabi, M., N. Alirezazadeh and H. T. Ahmadi. 1998. A Survey Of ^{222}Rn Concentrations in Domestic Water Supplies of Iran. *Health Physics*. 75: 417-421.
- [8] Yu, K., Z. Guan, Z. M. Stokes and E. Young. 1994. A Preliminary Study on the Radon Concentrations in Water in Hong Kong and the Associated Health Effects. *Applied Radiation and Isotopes*. 45: 809-810.
- [9] Garba, N. N., A. T. Ramli, H. T. Gabdo and M. S. Sanusi. 2013. Radiological Information of Kelantan-A Review. *Archives of Physics Research*. 4: 55-59.
- [10] Hamzah, Z., S. A. A. Rahman, S. M. Ahmad and S. Hamzah. 2012. Evaluation of Natural radioactivity in soil in district of Kuala Krai, Kelantan. *Malaysian Journal of Analytical Sciences*. 16: 335-345.
- [11] DGGs. 1982. Director General of Geological Survey. Map of Mineral Resources in Kelantan State, Malaysia 1st Edition. Ipoh, Malaysia.
- [12] Garba, N. N., A. T. Ramli, M. A. Saleh, M. S. Sanusi and H. T. Gabdo. 2014. Assessment of Terrestrial Gamma Radiation Dose Rate (TGRD) of Kelantan State, Malaysia: Relationship Between the Geological Formation and Soil Type to Radiation Dose Rate. *Journal of Radioanalytical and Nuclear Chemistry*. 302: 201-209.
- [13]. Paramanathan, S. 2000. Soils of Malaysia: Their Characteristics and Identification, ed. S. Paramanathan. Vol. 1. Kuala Lumpur, Malaysia: Academy of Sciences Malaysia.
- [14] Saleh, M. A., A. T. Ramli, Y. Alajerami, M. H. Abu Mhareb, A. S. Aliyu, H. T. Gabdo and N. N. Garba. 2014 Assessment of Radiological Health from Ambient Environment in the Muar District, Johor, Malaysia. *Radiation Physics and Chemistry*. 103: 243-252.
- [15] Peter, H. S. and D. H. Bruce. 1989. Radionuclides in Aquatic Environments. International Journal of Radiation Applications and Instrumentation. Part C. *Radiation Physics and Chemistry*. 34: 213-240.
- [16] Sarkar, B. 2002. *Heavy Metals in the Environment*. CRC Press.
- [17] Gabdo, H.T., A. T. Ramli, N. N. Garba, M. A. Saleh and M. S. Sanusi. 2015. Assessment of Natural Radionuclides in Rivers of Pahang state, Malaysia. In *3rd International Science Postgraduate Conference 2015 (ISPC2015) of Conference*. Ibnu Sina Institute for Fundamental Science Studies, Universiti Teknologi Malaysia 81310 Johor Bahru, Johor, Malaysia.
- [18] Khan, K., P. Akhter, S. Orfi, G. Malik and M. Tufail. 2003. Natural Radioactivity Levels in River, Stream and Drinking Water of the Northwestern Areas of Pakistan. *Journal of Radioanalytical and Nuclear Chemistry*. 256: 289-292.
- [19] Elles, M., A. Armstrong and S. Lee. 1997. Characterization and Solubility Measurements of Uranium-Contaminated Soils to Support Risk Assessment. *Health Physics*. 72: 716-726.

- [20] Ramli, A. T., M. A. AbdelWahab and M. S. Lee. 2001. Geological Influence on Terrestrial Gamma Ray Dose Rate in the Malaysian State of Johore. *Appl. Radiat. Isot.* 54: 327-333.
- [21] Saleh, M. A., A. T. Ramli, Y. Alajerami, A. S. Aliyu and N. A. B. Basri. 2013. Radiological study of Mersing District, Johor, Malaysia. *Radiation Physics and Chemistry.* 85: 107-117.